



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

## Journal of Threatened Taxa

Building evidence for conservation globally

[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

### NOTE

#### **NEW RECORDS OF *HELODERMA ALVAREZI* (WIEGMANN, 1829) (SAURIA: HELODERMATIDAE) ON THE COAST OF OAXACA AND INCREASES TO ITS DISTRIBUTION IN MEXICO**

Jesús García-Grajales, Rodrigo Arrazola Bohórquez, María Arely Penguilly Macías & Alejandra Buenrostro Silva

26 March 2020 | Vol. 12 | No. 4 | Pages: 15495–15498

DOI: [10.11609/jott.5691.12.4.15495-15498](https://doi.org/10.11609/jott.5691.12.4.15495-15498)



For Focus, Scope, Aims, Policies, and Guidelines visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>

For Policies against Scientific Misconduct, visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>

For reprints, contact [<ravi@threatenedtaxa.org>](mailto:ravi@threatenedtaxa.org)

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Member



Publisher & Host







## New records of *Heloderma alvarezii* (Wiegmann, 1829) (Sauria: Helodermatidae) on the coast of Oaxaca and increases to its distribution in Mexico

Jesús García-Grajales<sup>1</sup> , Rodrigo Arrazola Bohórquez<sup>2</sup> , María Arely Penguilly Macías<sup>3</sup>  &  
Alejandra Buenrostro Silva<sup>4</sup> 

<sup>1</sup> Instituto de Recursos, Universidad del Mar campus Puerto Escondido. Km. 2.5, Carr. Federal Puerto - Sola de Vega, Puerto Escondido 71980, Oaxaca, México.

<sup>2</sup> Universidad del Mar campus Huatulco. Ciudad Universitaria, Santa María Huatulco 70989, Oaxaca, México.

<sup>3</sup> Fondo Oaxaqueño para la Conservación de la Naturaleza. Puerto Escondido 71980, Oaxaca, México.

<sup>4</sup> Instituto de Industrias, Universidad del Mar campus Puerto Escondido. Km. 2.5, Carr. Federal Puerto - Sola de Vega, Puerto Escondido 71980, Oaxaca, México.

<sup>1</sup> [archosaurio@yahoo.com.mx](mailto:archosaurio@yahoo.com.mx), <sup>2</sup> [rodrigo.arrazolab@gmail.com](mailto:rodrigo.arrazolab@gmail.com), <sup>3</sup> [arely.penguilly@fondoax.org](mailto:arely.penguilly@fondoax.org),

<sup>4</sup> [sba\\_1575@yahoo.com.mx](mailto:sba_1575@yahoo.com.mx) (corresponding author)

In Mexico, the genus *Heloderma* is composed of five broad-ranging species: (i) the Gila Monster *H. suspectum* distributed in the Mohave Desert of extreme Nevada, southwestern Utah, extreme southeastern California and northwestern Arizona, and throughout the Sonoran Desert region in Arizona and Sonora, Mexico, as well as into the Chihuahuan Desert of southeastern Arizona and southwestern New Mexico (Campbell & Lamar 2004; Beck 2005); (ii) the Rio Fuerte Beaded Lizard *H. exasperatum* inhabits the foothills of the Sierra Madre Occidental, in the transition area between the drainage basins of Rio Mayo and Rio Fuerte and the Sonoran-Sinaloan subtropical dry forest in southern Sonora, extreme western Chihuahua, and northern Sinaloa (Campbell & Lamar 2004; Beck 2005); (iii) the Mexican Beaded Lizard *H. horridum* is found primarily in dry forest habitats from southern Sinaloa southward to Oaxaca and inland into the states of Mexico and Morelos (Bogert &

del Campo 1956; Campbell & Lamar 2004; Beck 2005); (iv) the Chiapan Beaded Lizard *H. alvarezii* inhabits dry forests in the Central Depression of central Chiapas and the Rio Lagartero Depression in extreme western Guatemala (Campbell & Lamar 2004; Beck 2005); and (v) the Guatemalan Beaded Lizard *H. charlesbogerti* that inhabits the Rio Motagua Valley in the Atlantic slope of eastern Guatemala (Campbell & Vannini 1988).

The beaded lizard is known colloquially as ‘escorpión’ and is well known to local inhabitants, yet its natural history is surrounded by mystery, notoriety and misconception (Reiserer et al. 2013; Domínguez-Vega et al. 2018), principally because these animals can pass up to 95% of their lives hidden in shelters underground, which makes it difficult to detect them. Moreover, they maintain a low population (Beck & Jennings 2003; Domínguez-Vega et al. 2018). Furthermore, it is the only group which bears venom-transmitting teeth

**Editor:** Anonymity requested.

**Date of publication:** 26 March 2020 (online & print)

**Citation:** Grajales, J.G., R.A. Bohórquez, M.A.P. Macías & A.B. Silva (2020). New records of *Heloderma alvarezii* (Wiegmann, 1829) (Sauria: Helodermatidae) on the coast of Oaxaca and increases to its distribution in Mexico. *Journal of Threatened Taxa* 12(4): 15495–15498. <https://doi.org/10.11609/jott.5691.12.4.15495-15498>


**Copyright:** © Grajales et al. 2020. Creative Commons Attribution 4.0 International License. JOTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

**Funding:** Universidad del Mar.

**Competing interests:** The authors declare no competing interests.

**Acknowledgements:** We give thanks to Universidad del Mar (UMAR) for the logistic support, and thank Alison Tai Rosewicz for your help with the language review. JGG thanks to Sistema Nacional de Investigadores (SNI) for the grant.





**Figure 1.** Location of photographic records of *Heloderma alvarezii* along the coast of Oaxaca; and historic published records (orange circles) in Chiapas State.

with a deep venom groove in the rostral carina (Fry et al. 2006). Although bites by the lizards are extremely rare, envenomation is known to produce severe local pain, dizziness, diaphoresis, vomiting, paresthesia, and hypotension (Ariano-Sánchez 2006). Consequently, the lizard is frequently slaughtered when encountered by humans (Beck 2005; Domínguez-Vega et al. 2017).

Herein, we provide three new records and a significant extension on the distribution of *H. alvarezii* about different encounters between humans while performing their activities and the species during its movement along the central coast of Oaxaca, Mexico, particularly in three different municipalities (Figure 1).

The first observation occurred on 04 July 2017, when a lizard was held in captivity in the town of Santa Rosa de Lima, municipality of Villa de Tututepec de Melchor Ocampo. This animal was used as an exhibition specimen to decrease the fear of this species in the town. We were prohibited from measuring it, weighing it or determining its sex; however, to the naked eye this individual presented good body condition (Image 1a). According to the people in charge of the lizard, it came from the upper zone, close to the municipality.

The second observation occurred on 10 October 2019 at 17.15 h. An adult female *H. alvarezii* was found and captured by people who were cleaning a field in a locality known as ‘El Porvernir’ in the Municipality of Santa María Colotepec, Oaxaca. Due to their fear of the

creature, people caught the lizard in order to slaughter it, but then they reflected on the situation and reported it to the Universidad del Mar in the city of Puerto Escondido. The individual was measured (total length: 670mm), photographed (Image 1b) and then traslocated and released into the Natural Protected Area El Gavilán in San Francisco Cozoaltepec, municipality of Santa María Tonameca, Oaxaca, to ensure that the individual would not be harmed.

Most recently, the third observation occurred on 03 November 2019 at 18.20h, between the boundaries of the Universidad del Mar campus Huatulco and Huatulco National Park, in the Municipality of Santa María Huatulco. A single juvenile was found walking near a road and was later photographed on a tree (Image 2c). It was not manipulated in order to avoid causing it unnecessary stress.

The identification of these lizards was possible because this taxon is unique among the beaded lizards in that it undergoes an ontogenetic increase in melanism (Bogert & del Campo 1956; Beck 2005). Also distinctive is that yellow banding on the tail, a characteristic typical of the other species of beaded lizards, is essentially absent in adults of *H. alvarezii* (Bogert & del Campo 1956; Beck 2005). Although Bogert & del Campo (1956) and Beck (2005) explain that black individuals are uncommon, there is recent photographic evidence of black individuals on the coast of Oaxaca. Neonates and



**Image 1.** *Heloderma alvarezii* in: a—Villa de Tututepec de Melchor Ocampo municipality | b—Santa María Colotepec municipality | c—Santa María Huatulco municipality.

juveniles often are distinctly marked with yellow spots and bands on the tail, but the color pattern of adults gradually transforms to an almost uniform dark brown or gray coloration.

*Heloderma alvarezii* was described as inhabiting dry tropical forest in the Central Depression (Río Grijalva Depression) of central Chiapas and the Río Lagartero Depression in extreme western Guatemala (Campbell & Lamar 2004; Beck 2005; Köhler 2008; Johnson et al. 2010; Wilson et al. 2010). Nonetheless, Álvarez del Toro (1983 in Campbell & Vannini 1988) indicated the probability of sympatry areas between *H. horridum* and *H. alvarezii* in the region between the Isthmus of Tehuantepec (Oaxaca) and Cintalapa (Chiapas); however, up to this point, the literature on integration between these two species is inconclusive (Reiserer et al. 2013), and therefore, the most recent herpetofauna list in Oaxaca does not include *H. alvarezii*'s presence in the State. Sánchez de la Vega et al. (2012) provided, however, the photographic record of *H. horridum* in the Municipality of San Pedro Mixtepec, and previous records were made of this

species in Jamiltepec and the Isthmus of Tehuantepec (Bogert & del Campo 1956).

Helodermatid lizards are listed as “threatened” under Mexican law (NOM-059-SEMARNAT-2010, SEMARNAT 2010) and assessed as ‘Vulnerable’ by the Red List of International Union for Conservation of Nature (IUCN). The main threat to beaded lizards populations is primarily the deforestation of dry tropical forests for agriculture, cattle ranching, and the burgeoning human population (Janzen 1988; Myer et al. 2000; Williams-Linera & Lorea 2009; Pennington et al. 2006; Domínguez-Vega et al. 2012), as well as, an escalation of droughts and fires (Beck 2005; Miles et al. 2006). Additionally, anthropogenic pressure is a threat.

These three photographic records represent the most northeastern (NE) geographic records of *H. alvarezii*, and it increases the distribution of the species approximately 359km for the record 3, 405km for record 2, and 470km for record 1 to the northeast from the previously closest reported localities of Cintalapa and Ocozocoautla, Chiapas. It is necessary to gather more records on the

distribution of this Mexican species to amend the lack of information about this. In conclusion, our records of this species increase the number of reptiles present in Oaxaca to 448 species, if we considered the recent contributions made by Carbajal-Marquez et al. (2020).

## References

- Ariano-Sánchez, D. (2006).** The Guatemalan beaded lizard: endangered inhabitant of a unique ecosystem. *Iguana* 13: 179–183.
- Beck, D.D. (2005).** *Biology of Gila monsters and Beaded lizards*. University of California Press, Berkeley, California, USA, 247pp.
- Beck, D.D. & R. Jennings (2003).** Habitat use by Gila monsters: the importance of shelters. *Herpetological Monographs* 17: 111–129.
- Bogert, C.M. & R.M. del Campo (1956).** The Gila monster and its allies: the relationships, habits, and behaviour of the lizards of the family Helodermatidae. *Bulletin of the American Museum of Natural History* 109: 1–238.
- Campbell, J.A. & W.W. Lamar (2004).** The venomous reptiles of the Western Hemisphere (2 volumes). Comstock Publishing Associates, Cornell University Press, Ithaca, New York, USA, 528pp.
- Campbell, J.A. & J.P. Vannini (1988).** A new subspecies of beaded lizard, *Heloderma horridum*, from the Motagua Valley of Guatemala. *Journal of Herpetology* 22: 457–468.
- Domínguez-Vega, H., O. Monroy-Vilchis, C.J. Balderas-Valdivia, C.M. Gienger & D. Ariano-Sánchez (2012).** Predicting the potential distribution of the beaded lizard and identification of priority areas for conservation. *Journal of Nature Conservation* 20: 247–253. <https://doi.org/10.1016/j.jnc.2012.04.003>
- Domínguez-Vega, H., O. Monroy-Vilchis, J. Manjarrez & C.J. Balderas-Valdivia (2017).** Aversive hunting and sight frequency ecology of Beaded lizards (Squamata: Helodermatidae). *Perspective in Ecology and Conservation* 15: 47–51. <https://doi.org/10.1016/j.pecon.2016.11.003>
- Domínguez-Vega, H., C.J. Balderas-Valdivia, J. Manjarrez & O. Monroy-Vilchis (2018).** Conociendo al lagarto escorpión: leyendas, realidad y potencial de una rareza biológica. *Ciencia ergo-sum* 25(2): 1–8. <https://doi.org/10.30878/ces.v25n2a10>
- Fry, B.G., N. Vidal, J.A. Norman, F.J. Vonk, H. Scheib, S.F.R. Ramjan, S. Kuruppu, K. Fung, S.B. Hedges, M. Richardson, W.C. Hodgson, V. Ignjatovic, R. Summerhayes & E. Kochva (2006).** Early evolution of the venom system in lizards and snakes. *Nature* 439: 584–588.
- Janzen, D.H. (1988).** Tropical dry forest: the most endangered major tropical ecosystem. pp. 130–137. In: Wilson, E.O. (ed.). *Biodiversity*. National Academy Press, Washington, D. C., USA, 538pp.
- Johnson, J.D., V. Mata-Silva & L.D. Wilson (2010).** Geographic distribution and conservation of the herpetofauna of southeastern Mexico, pp. 322–369 In: Wilson, L.D., J. Townsend & J.D. Johnson (eds.). *Conservation of mesoamerican amphibians and reptiles* Eagle Mountain Publishing, LC., Eagle Mountain, Utah, United States.
- Köhler, G. (2008)** *Reptiles of Central America*. Herpeton, Offenbach, Germany, xvii+812pp.
- Miles, L., A.C. Newton, D.S. DeFries, C. Ravilious, I. May, S. Blyth, V. Kapos & J.E. Gordon (2006).** A global overview of the conservation status of tropical dry forests. *Journal of Biogeography* 33: 491–505.
- Pennington, R.T., G.P. Lewis & J.A. Ratter (2006).** *Neotropical savannas and seasonally dry forests: Plant diversity, biogeography, and Conservation*. CRC Press, Boca Raton, Florida, USA, 508pp.
- Reiserer, R.S., G.W.S. Schuett & D. Beck (2013).** Taxonomic reassessment and conservation status of the beaded lizard, *Heloderma horridum* (Squamata: Helodermatidae). *Amphibian & Reptile Conservation* 7(1): 74–96.
- Sánchez de la Vega, G., A.B. Silva, J.G. Grajales & V.M. Silva (2012).** Geographic distribution. *Heloderma horridum* (Mexican beaded lizard). Mexico: Oaxaca. *Herpetological Review* 43: 102.
- SEMARNAT (2010).** Norma Oficial Mexicana NOM-059-SEMARNAT-2010, que determina las especies de flora y fauna silvestres, terrestres y acuáticas, endémicas, amenazadas, en peligro de extinción y sujetas a protección especial. Órgano del Gobierno Constitucional de los Estados, Gobierno Federal, México.
- Williams-Linera, G. & F. Lorea (2009).** Tree species diversity driven by environmental and anthropogenic factors in tropical dry forest fragments in central Veracruz, Mexico. *Biodiversity and Conservation* 18: 3269–3293.







PLATINUM  
OPEN ACCESS



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

March 2020 | Vol. 12 | No. 4 | Pages: 15407–15534

Date of Publication: 26 March 2020 (Online & Print)

DOI: 10.11609/jott.2020.12.4.15407-15534

[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

### Conservation Application

**Do wildlife crimes against less charismatic species go unnoticed? A case study of Golden Jackal *Canis aureus* Linnaeus, 1758 poaching and trade in India**

– Malaika Mathew Chawla, Arjun Srivathsa, Priya Singh, Iravatee Majgaonkar, Sushma Sharma, Girish Punjabi & Aditya Banerjee, Pp. 15407–15413

### Review

**Hazards of wind turbines on avifauna - a preliminary appraisal within the Indian context**

– Himika Deb, Tanmay Sanyal, Anilava Kaviraj & Subrata Saha, Pp. 15414–15425

### Communications

**Analysis of stereotypic behaviour and enhanced management in captive Northern Giraffe *Giraffa camelopardalis* housed at Zoological Garden Alipore, Kolkata**

– Tushar Pramod Kulkarni, Pp. 15426–15435

**A new species of shieldtail snake (Reptilia: Squamata: Uropeltidae) from Kolli Hill complex, southern Eastern Ghats, peninsular India**

– S.R. Ganesh & N.S. Achyuthan, Pp. 15436–15442

**The insect fauna of Tenompok Forest Reserve in Sabah, Malaysia**

– Arthur Y.C. Chung, Vivianny Paul & Steven Bosuang, Pp. 15443–15459

**Tiger beetles (Coleoptera: Cicindelinae) of Davao Region, Mindanao, Philippines**

– Milton Norman Medina, Analyn Cabras, Harlene Ramillano & Reagan Joseph Villanueva, Pp. 15460–15467

**An assessment of the conservation status of a presumed extinct tree species *Wendlandia angustifolia* Wight ex. Hook.f. in southern Western Ghats, India**

– Chellam Muthumperumal, Paramasivam Balasubramanian & Ladan Rasingam, Pp. 15468–15474

### Short Communications

**Additional morphological notes on the male of *Icius alboterminus* (Caleb, 2014) (Aranei: Salticidae) with new distribution records from India**

– Dhruv A. Prajapati & R.D. Kamboj, Pp. 15475–15480

**Three moss families (Bryopsida: Calymperaceae, Hyopterygiaceae, & Pterobryaceae): new distribution records to bryoflora of Andhra Pradesh, India**

– Ananthaneni Sreenath, Midigesi Anil Kumar, Paradesi Anjaneyulu & Boyina Ravi Prasad Rao, Pp. 15481–15488

### Notes

**Mating behavior of the Yellow-throated Marten *Martes flavigula* (Mammalia: Carnivora: Mustelidae)**

– Abinash Parida, Meesala Krishna Murthy & G.S. Solanki, Pp. 15489–15492

**New to Myanmar: the Rosy Starling *Pastor roseus* (Aves: Passeriformes: Sturnidae) in the Hkakabo Razi Landscape**

– Sai Sein Lin Oo, Myint Kyaw, Nay Myo Hlaing & Swen C. Renner, Pp. 15493–15494

**New records of *Heloderma alvarezii* (Wiegmann, 1829) (Sauria: Helodermatidae) on the coast of Oaxaca and increases to its distribution in Mexico**

– Jesús García-Grajales, Rodrigo Arrazola Bohórquez, María Arely Penguilly Macías & Alejandra Buenrostro Silva, Pp. 15495–15498

**Description of a new subspecies of the genus *Microcerotermes* Silvestri, 1901 (Amitermitinae: Termitidae: Isoptera) and the first record of another termite species from Meghalaya, India**

– Khirod Sankar Das & Sudipta Choudhury, Pp. 15499–15502

**A new record of the hoverfly genus *Dasysyrphus* Enderlein, 1938 (Insecta: Diptera: Syrphidae) from India**

– Jayita Sengupta, Atanu Naskar, Aniruddha Maity, Panchanan Parui, Sumit Homchaudhuri & Dhriti Banerjee, Pp. 15503–15506

**First record of Banded Lineblue *Prosotas aluta* Druce, 1873 (Insecta: Lepidoptera: Lycaenidae) from Bangladesh**

– Rajib Dey, Ibrahim Khalil Al Haidar, Sajib Rudra & M. Rafiqul Islam, Pp. 15507–15509

**Notes on *Ptilomera agriodes* (Hemiptera: Heteroptera: Gerridae) from Eastern Ghats, India**

– J. Deepa, A. Narahari, M. Karuthapandi, S. Jadhav & C. Shiva Shankar, Pp. 15510–15513

***Didymocarpus bhutanicus* W.T. Wang (Gesneriaceae): a new addition to the herbs of India**

– Subhajit Lahiri, Sudhansu Sekhar Dash, Monalisa Das & Bipin Kumar Sinha, Pp. 15514–15517

**Rediscovery of *Epilobium trichophyllum* Hausskn.: a rare and endemic plant from Sikkim Himalaya, India**

– David L. Biate & Dinesh K. Agrawala, Pp. 15518–15521

**Additions of woody climbers (Lianas) to the flora of Manipur, India**

– Longjam Malemnganbee Chanu & Debjyoti Bhattacharyya, Pp. 15522–15529

**Molecular characterization of stinkhorn fungus *Aseroë coccinea* Imazeki et Yoshimi ex Kasuya 2007 (Basidiomycota: Agaricomycetes: Phallales) from India**

– Vivek Bobade & Neelesh Dahanukar, Pp. 15530–15534

Member



Publisher & Host

